

# NAVAL POSTGRADUATE SCHOOL Monterey, California



## ASYMPTOTIC PROPERTIES OF A SENSOR ALLOCATION MODEL

by

Donald P. Gaver  
Patricia A. Jacobs  
Mark Youngren

May 1995

Approved for public release; distribution is unlimited

Prepared for: Institute for Joint Warfare Analysis, Monterey, CA 93943-5000  
and  
Force Structure, Resources and Assessment Directorate (J8),  
The Joint Staff, Washington, DC 20350-5000



NAVAL POSTGRADUATE SCHOOL  
MONTEREY, CA 93943-5000

Rear Admiral T. A. Mercer  
Superintendent

Harrison Shull  
Provost

This report was prepared for and funded by the Institute for Joint Warfare Analysis, Monterey, CA, and the Force Structure, Resources, and Assessment Directorate (J8), The Joint Staff, Washington DC.

Reproduction of all or part of this report is authorized.

This report was prepared by:



**REPORT DOCUMENTATION PAGE**Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)

2. REPORT DATE

May 1995

3. REPORT TYPE AND DATES COVERED

Technical

4. TITLE AND SUBTITLE

Asymptotic Properties of a Sensor Allocation Model

5. FUNDING NUMBERS

RLGPD

6. AUTHOR(S)

Donald P. Gaver, Patricia A. Jacobs and Mark Youngren

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Naval Postgraduate School  
Monterey, CA 939438. PERFORMING ORGANIZATION  
REPORT NUMBER

NPS-OR-95-002

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Institute for Joint Warfare Analysis, Monterey, CA 93943-5000  
Force Structure, Resources and Assessment Directorate (J8),  
The Joint Staff, Washington DC, 20350-500010. SPONSORING / MONITORING  
AGENCY REPORT NUMBER

11. SUPPLEMENTARY NOTES

12a. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release; distribution is unlimited.

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)

A dynamic adaptive protocol for allocating sensor assets to locations where most opponents' assets appear is described, analyzed, and illustrated.

14. SUBJECT TERMS

adaptive allocation; estimation of the number of binomial trials;  
normal approximations

15. NUMBER OF PAGES

36

16. PRICE CODE

17. SECURITY CLASSIFICATION  
OF REPORT

Unclassified

18. SECURITY CLASSIFICATION  
OF THIS PAGE

Unclassified

19. SECURITY CLASSIFICATION  
OF ABSTRACT

Unclassified

20. LIMITATION OF ABSTRACT

UL





# Asymptotic Properties of a Sensor Allocation Model

D. P. Gaver

P. A. Jacobs

M. Youngren

## 0. Introduction and Summary

Sensor resources are to be allocated among  $I$  non-overlapping but possibly contiguous geographical locations called *nodes*. Each sensor look at a node results in an observation, with error, of the number of *units* that are at the node. In military applications *units* may be individual assets such as tanks or ships, or possibly small groups organized as platoons or companies and in geographical proximity; in ecological applications, they might be animals, singly or in groups, or particular vegetation types. The observations are then to be used to estimate the numbers of units at each node. The problem is to allocate sensor resources so as to minimize a measure of the error of the estimates, and of the estimate of the sum of the numbers at all nodes. It is assumed that the units remain on the nodes and do not migrate, although for some sensor types the units must be in motion on the node at least some of the time in order for detection to take place.

The objective of this paper is to propose and study *adaptive allocation* of sensor effort in such a way as to focus the sensor's attention sequentially and purposefully on nodes so as to pay most attention to those nodes about which the greatest uncertainty, or interest, currently prevails. This problem has features in common with adaptive bin-packing problems (cf. Gaver et al., 1995) and adaptive allocation of customers to servers (cf. Gaver et al., 1993).

Numerical examples show that a properly selected sequentially adaptive rule will provide estimates of improved precision.

A related investigation that has recently come to our attention is that by Thompson and Seber (1994). Their procedures can also be analyzed using the methodology proposed in this paper.

## 1. The Model

There are  $I$  nodes. Assume that node  $i$  contains  $r_i$  units. Sensor resources are allocated to one of the  $I$  nodes at times that occur according to a Poisson process with rate  $\lambda$ . The sensor resource is allocated to node  $i$  with probability  $\alpha_i$  which is tailored to depend on past allocations in a purposeful way. Let  $Z_n(i)$  be the  $n^{\text{th}}$  observation of node  $i$ ; a simple model is that  $\{Z_n(i); n = 1, 2, \dots\}$  are iid with binomial distribution with  $r_i$  trials and known probability of success  $p_i$  which is the probability of unit detection at node  $i$ . We assume for illustration that the number of units on a node does not change, although an adaptive scheme of the type proposed should effectively follow changes in node population.

Let  $N_i(t)$  be the number of times in  $[0, t]$  that node  $i$  is observed by a sensor.

Let

$$V(i; t) = Z_1(i) + \dots + Z_{N_i(t)}(i)$$

be the sum of all the observations during  $[0, t]$  for node  $i$ .

Under some conditions  $V(i; t)$  has a binomial distribution with  $N_i(t)r_i$  trials and probability of success  $p_i$ , given  $N_i(t)$ , although this assumption may not be especially accurate in general. Assuming it to be adequate for the moment, an estimate for the number of units at node  $i$  based on observations made during  $[0, t]$  is

$$\hat{r}_i(t) = \frac{V(i; t)}{N_i(t)p_i} \quad (1.1)$$



with

$$\text{Var}[\hat{r}_i(t)] = \frac{r_i N_i(t) p_i (1 - p_i)}{N_i(t)^2 p_i^2} \quad (1.2)$$

$$= \frac{r_i (1 - p_i)}{N_i(t) p_i} \approx \frac{\hat{r}_i(t) (1 - p_i)}{N_i(t) p_i} = \frac{V(i; t) (1 - p_i)}{N_i(t)^2 p_i^2}. \quad (1.3)$$

There has been much statistical attention paid to estimating the number of trials in a binomial distribution, given probability of success; for recent discussion see Hall (1994). Here we use the simplest such estimator.

Now consider the following *adaptive allocation rule*. A sensor resource arriving at time  $t$  is allocated to node  $i$  with probability

$$\alpha_i(N_i(t), V_i(t); N(t), V(t)) = \frac{h_i(N_i(t), V_i(t))}{\sum_{j=1}^I h_j(N_j(t), V_j(t))} \quad (1.4)$$

where the  $h_i$  are strictly positive sufficiently smooth functions such that  $h_j(c^p x, c^p y) = c^p h_j(x, y)$ .

One possible form for  $h_i$  is

$$h_i(x, v) = \left[ \frac{a_i}{x} + \frac{v(1 - p_i)}{x^2 p_i^2} \right]^\gamma \quad \text{for } \gamma > 0 \text{ and } a_i > 0; \quad (1.5)$$

if it is assumed that given  $N_i(t)$ ,  $V_i(t)$  is binomially distributed with mean  $N_i(t)p_i$ , then from (1.3) it follows that for this function  $h_i$ , sensor resources tend to be allocated to those nodes for which the variance, and hence its square root, the standard error of the estimated number of units, is the largest; the probability of allocation to the most uncertain node (by this measure) increases rapidly, approaching unity as  $\gamma$  increases. This tends to bring down that standard error quickly, and to equate standard errors of the estimates across nodes. Clearly, alternative measures of overall sensor performance are feasible, and possibly desirable, such as ones that endeavor to equalize fractional or percent error on

nodes, or ones that also respond to an independent measure of importance of the units on a node. Additionally, node contents may be of various types, which can be considered. Thus, the present discussion is of an illustration of an adaptive allocation scheme.

The purposeful allocation of (1.4) introduces dependence between  $\{N_i(t); t \geq 0\}$  and  $\{V_i(t); t \geq 0\}$ . Asymptotic results for the means  $E[N_i(t)]$  and  $E[V_i(t)]$  as the rate of the Poisson process  $\lambda \rightarrow \infty$  are obtained in Section 2. It is shown that the purposeful allocation estimate  $\hat{r}_i(t)$  is asymptotically unbiased. Section 3 presents results for the asymptotic means for specific form of function  $h_i$  (1.5). Section 4 discusses asymptotic results for the second moments of  $\{N_i(t); t \geq 0\}$  and  $\{V_i(t); t \geq 0\}$  and presents approximate expressions for the  $Var[\hat{r}_i(t)]$  and  $Var\left[\sum_i \hat{r}_i(t)\right]$ . Section 5 describes a simpler Poisson approximation. Section 6 presents results from simulation experiments.

## 2. First-Moment Calculations and Asymptotics

Note that for  $h > 0$

$$E[N_i(t+h)|N(t), V(t)] = N_i(t) + \lambda \alpha_i(N, V)h. \quad (2.1)$$

Thus

$$E[N_i(t+h)] = E[N_i(t)] + \lambda E[\alpha_i(N, V)h]. \quad (2.2)$$

Assuming derivatives exist, we have

$$\frac{d}{dt} E[N_i(t)] = \lambda E[\alpha_i(N(t), V(t))]. \quad (2.3)$$

Similarly,

$$E[V_i(t+h)|N(t), V(t)] = V_i(t) + \lambda h \alpha_i(N, V) r_i p_i. \quad (2.4)$$

Thus,

$$E[V_i(t+h)] = E[V_i(t)] + \lambda r_i p_i E[\alpha_i(N, V)h]. \quad (2.5)$$

Assuming derivatives exist, we have

$$\frac{d}{dt} E[V_i(t)] = r_i p_i \lambda E[\alpha_i(N, V)]. \quad (2.6)$$

Assume  $\alpha_i$  is of the form (1.4) where  $h_i$  is sufficiently smooth with  $h_i(c^p x, c^p y) = c^p h_i(x, y)$  for constants  $c$  and  $h_i(x, y) > 0$  for all  $x, y > 0$ . We have

$$\begin{aligned} \alpha_i(N(t), V(t)) &= \frac{h_i(N_i(t), V_i(t))}{\sum_j h_j(N_j(t), V_j(t))} \\ &= \frac{h_i(N_i(t)/\lambda, V_i(t)/\lambda)}{\sum_j h_j(N_j(t)/\lambda, V_j(t)/\lambda)}. \end{aligned} \quad (2.7)$$

Assume  $\lim_{\lambda \rightarrow \infty} N_i(t)/\lambda = m_i(t)$ ,  $\lim_{\lambda \rightarrow \infty} V_i(t)/\lambda = v_i(t)$ ,  $\lim_{\lambda \rightarrow \infty} \frac{d}{dt} E[N_i(t)/\lambda] = \frac{d}{dt} m_i(t)$ , and  $\lim_{\lambda \rightarrow \infty} \frac{d}{dt} E[V_i(t)/\lambda] = \frac{d}{dt} v_i(t)$ .

Dividing both sides of (2.3) and (2.6) by  $\lambda$  and letting  $\lambda \rightarrow \infty$ , the bounded convergence theorem yields

$$\frac{d}{dt} m_i(t) = \frac{h_i(m_i(t), v_i(t))}{\sum_j h_j(m_j(t), v_j(t))}. \quad (2.8)$$

and

$$\frac{d}{dt} v_i(t) = r_i p_i \frac{h_i(m_i(t), v_i(t))}{\sum_j h_j(m_j(t), v_j(t))} \quad (2.9)$$

Thus,

$$v_i(t) = r_i p_i m_i(t). \quad (2.10)$$

### 3. Purposeful Allocation

The equations for  $m_i(t)$ , (2.8), can be solved for special functions  $h_i$ . For example, assume

$$h_i(x, y) = \left[ \frac{a_i}{x} + \frac{y(1-p_i)}{x^2 p_i^2} \right]^\gamma \quad \text{for } \gamma \geq 0 \text{ and } a_i > 0. \quad (3.1)$$

Since

$$v_i(t) = m_i(t) r_i p_i, \quad (3.2)$$

$$h_i(m_i(t), v_i(t)) = \left[ \frac{a_i}{m_i(t)} + \frac{r_i(1-p_i)}{m_i(t) p_i} \right]^\gamma.$$

In this case (2.8) can be rewritten as

$$\frac{\frac{d}{dt} m_i(t)}{\frac{d}{dt} m_j(t)} = \left[ \frac{a_i + r_i c_i}{a_j + r_j c_j} \frac{m_j(t)}{m_i(t)} \right]^\gamma \quad (3.3)$$

where  $c_i = (1 - p_i)/p_i$ .

The functions

$$m_i(t) = K(a_i + r_i c_i)^{\gamma/(\gamma+1)} t \quad (3.4)$$

with

$$K = \left[ \sum_{i=1}^I (a_i + r_i c_i)^{\gamma/(\gamma+1)} \right]^{-1} \quad (3.5)$$

are a solution to (2.8).

Thus, in the limit as  $t \rightarrow \infty$ , the probability that a sensor resource at time  $t$  looks at node  $i$  is

$$\alpha_i = \frac{(a_i + r_i c_i)^{\gamma/(\gamma+1)}}{\sum_j (a_j + r_j c_j)^{\gamma/(\gamma+1)}}. \quad (3.6)$$

If  $\gamma = 0$ , then

$$\alpha_i = 1/I \quad (3.7)$$

and the allocation is equally likely.

If  $\gamma \rightarrow \infty$ , then

$$\alpha_i = \frac{a_i + r_i c_i}{\sum_j a_j + r_j c_j}. \quad (3.8)$$

If, further, the probability of detection on node  $i$ ,  $p_i$ , is constant for all the nodes, then the probabilistic allocation for  $\gamma \rightarrow \infty$  is roughly proportional to the number of units on the node.

#### 4. Scaling and Approximate Variances of Estimators

Let

$$X_j(t) = \frac{N_j(t) - \lambda m_j(t)}{\sqrt{\lambda}} \quad (4.1)$$

$$Y_j(t) = \frac{V_j(t) - \lambda v_j(t)}{\sqrt{\lambda}} \quad (4.2)$$

In Appendix A, moment generating functions are used to show the asymptotic normality of  $\{(X_j(t), Y_j(t)); t \geq 0\}$  as  $\lambda \rightarrow \infty$ . Further, differential equations for their second moments are obtained.

Rewriting

$$N_j(t) = \lambda m_j(t) + \sqrt{\lambda} X_j(t) \quad (4.3)$$

$$V_j(t) = \lambda v_j(t) + \sqrt{\lambda} Y_j(t) \quad (4.4)$$

An approximate variance of the estimator of the number of units on node  $i$ ,

$$\hat{r}_i(t) = \frac{V_i(t)}{N_i(t)} \frac{1}{p_i} \quad (4.5)$$

can be computed using the "delta method" as follows; cf. Bickel and Doksum (1977). To begin,

$$\text{Var}[\hat{r}_i(t)] = \frac{1}{p_i^2} \text{Var}\left[\frac{V_i(t)}{N_i(t)}\right]. \quad (4.6)$$

A Taylor expansion yields

$$\begin{aligned} \frac{V_i(t)}{N_i(t)} &= \frac{v_i(t) + Y_i(t)/\sqrt{\lambda}}{m_i(t) + X_i(t)/\sqrt{\lambda}} \\ &= \frac{v_i(t)}{m_i(t)} + \frac{1}{m_i(t)} [Y_i(t)/\sqrt{\lambda}] - \frac{v_i(t)}{m_i(t)^2} (X_i(t)/\sqrt{\lambda}) \\ &\quad + O\left(\frac{1}{\lambda}\right). \end{aligned} \quad (4.7)$$

Thus,

$$\text{Var}\left[\frac{V_i(t)}{N_i(t)}\right] \approx E\left[\left(\frac{V_i(t)}{N_i(t)} - \frac{v_i(t)}{m_i(t)}\right)^2\right] \quad (4.8)$$

$$\begin{aligned} &\approx \frac{1}{\lambda m_i(t)^2} \left\{ E[Y_i^2(t)] + \left[\frac{v_i(t)}{m_i(t)}\right]^2 E[X_i^2(t)] - 2\frac{v_i(t)}{m_i(t)} E[Y_i(t)X_i(t)] \right\} \\ &= \frac{1}{\lambda m_i(t)^2} \left\{ E[Y_i^2(t)] + (r_i p_i)^2 E[X_i^2(t)] - 2r_i p_i E[X_i(t)Y_i(t)] \right\}. \end{aligned} \quad (4.9)$$

Hence, an approximate variance of the estimate of the number of units on node  $i$

$$\text{Var}[\hat{r}_i(t)] \approx \frac{1}{p_i^2} \frac{1}{\lambda m_i(t)^2} \left\{ E[Y_i^2(t)] + (r_i p_i)^2 E[X_i^2(t)] - 2r_i p_i E[X_i(t)Y_i(t)] \right\}. \quad (4.10)$$



To approximate the variance of the sum of the estimates, a Taylor expansion yields

$$\begin{aligned}
 E\left[\frac{V_i(t)}{N_i(t)}\frac{V_j(t)}{N_j(t)}\right] &= E\left[\frac{v_i(t)+Y_i(t)/\sqrt{\lambda}}{m_i(t)+X_i(t)/\sqrt{\lambda}}\frac{v_j(t)+Y_j(t)/\sqrt{\lambda}}{m_j(t)+X_j(t)/\sqrt{\lambda}}\right] \\
 &= \frac{v_i(t)v_j(t)}{m_i(t)m_j(t)} + \frac{1}{\lambda}\frac{1}{m_i(t)m_j(t)}E[Y_i(t)Y_j(t)] \\
 &\quad - \frac{v_i(t)v_i(t)}{\lambda[m_i(t)m_i(t)]^2}E[X_i(t)X_j(t)] + o\left(\frac{1}{\lambda}\right)
 \end{aligned} \tag{4.11}$$

for  $i \neq j$ .

Thus,

$$\begin{aligned}
 \text{Cov}(\hat{r}_i(t), \hat{r}_j(t)) &= \frac{1}{p_i p_j} \text{Cov}\left(\frac{V_i(t)}{N_i(t)}, \frac{V_j(t)}{N_j(t)}\right) \\
 &\approx \frac{1}{\lambda} \frac{1}{p_i m_i(t)} \frac{1}{p_j m_j(t)} \left\{ E[Y_i(t)Y_j(t)] - r_i p_i r_j p_j E[X_i(t)X_j(t)] \right\}.
 \end{aligned} \tag{4.12}$$

Expressions (4.10) and (4.12) can be used to approximate  $\text{Var}\left[\sum_i \hat{r}_i(t)\right]$ .

## 5. A Simpler Poisson Approximation

Assume the probability of allocation  $\alpha_i$  is independent of  $(V_i(t), N_i(t))$ ; then the number of looks at node  $i$  is a Poisson process with rate  $\lambda\alpha_i$  independent of the other nodes. Further

$$E[N_i(t)] = 1 + \alpha_i t \tag{5.1}$$

$$E[V_i(t)] = r_i p_i [1 + \alpha_i t] \tag{5.2}$$

$$\text{Var}[N_i(t)] = \alpha_i t \tag{5.3}$$

$$\text{Var}[V_i(t)] = r_i p_i (1 - p_i) + \alpha_i t \left[ r_i p_i (1 - p_i) + (r_i p_i)^2 \right] \tag{5.4}$$

$$\text{Cov}[N_i(t), V_i(t)] = r_i p_i \alpha_i t \quad (5.5)$$

where we assume that at time 0 each node is looked at once.

In this case (4.10) becomes

$$\text{Var}[\hat{r}_i(t)] = \frac{1}{p_i^2} \frac{r_i p_i (1 - p_i)}{1 + \alpha_i t}. \quad (5.6)$$

and the estimators  $\hat{r}_i(t)$  are independent.

Assume purposeful allocation is adapted with function  $h_i$  as in (3.1). A simple approximation to  $\text{Var}\left[\sum_i \hat{r}_i(t)\right]$  can be obtained by neglecting all covariances and assuming the number of looks at node  $i$ ,  $\{N_i(t); t \geq 0\}$ , is a Poisson process with rate  $\lambda \alpha_i$  where  $\alpha_i$  is determined by (3.6). In this case

$$\text{Var}\left[\sum_i \hat{r}_i(t)\right] \approx \sum_i \frac{1}{p_i} \frac{r_i (1 - p_i)}{(1 + \alpha_i t)}. \quad (5.7)$$

## 6. Numerical Examples

Suppose there are 3 nodes with  $r_j$  units on node  $j$  with  $r_1 = 49$ ,  $r_2 = 25$ , and  $r_3 = 16$ . The probabilities of detecting a unit on node  $j$ ,  $p_j$ , are  $p_1 = 1/11$ ,  $p_2 = 0.5$ , and  $p_3 = 10/11$ .

The variance of the estimate of the sum of the units on all the nodes under purposeful allocation with  $h_i$  as in (3.1) was studied using simulation for  $\gamma = 0, 1, 10$ . Each replication of the simulation begins with one observation at each node. The times of arrival of a Poisson process with rate 1 are then simulated. A node for observation at time  $t$  is randomly chosen using probabilities

$$\alpha_i(t) = K(t) \left[ \frac{a_i}{N_i(t)} + \frac{V_i(t)(1 - p_i)}{N_i(t)^2 p_i^2} \right]$$

for  $i = 1, 2, 3$  with  $K(t)$  the normalizing constant. A binomial observation is generated for the node chosen. The simulation has 500 replications.

Table 1 records the sample mean and square root of the sample variance of the sum of the estimates  $\sum_{i=1}^3 \hat{r}_i(t)$  for  $t = 5, 10, 20, 50$  where

$$\hat{r}_i(t) = \frac{V_i(t)}{N_i(t)p_i}.$$

Also shown are the sample means of the square root of the sum of estimated approximate variances (1.3)

$$\hat{\sigma}_i^2(t) = \frac{V_i(t)(1-p_i)}{N_i(t)^2 p_i^2}.$$

**Table 1**  
**Estimate of Total Number of Units on All Nodes**  
 $r_1 = 49, r_2 = 25, r_3 = 16; p_1 = 1/11, p_2 = 0.5, p_3 = 10/11$

		Simulation			Approximation	
		$\sum_i \hat{r}_i(t)$		$\sqrt{\sum_i \sigma_i^2(t)}$	Square Root of Differential Equation Approx Variance	Square Root of Poisson Approx Variance
Time	$\gamma$	Sample Mean	Square Root of Sample Variance	Sample Mean		
5	0	89.8	18.0	15.0	13.9	13.9
	1	88.1	15.1	10.7	12.3	10.7
	10	89.3	15.2	10.9	12.5	10.4
10	0	90.3	12.4	11.9	10.9	10.9
	1	89.8	9.2	8.2	9.1	8.1
	10	90.1	10.2	8.6	10.2	8.0
20	0	90.0	8.9	8.6	8.2	8.2
	1	89.5	7.1	6.1	6.7	6.0
	10	89.5	8.3	6.5	8.0	6.1
50	0	89.9	5.8	5.5	5.4	5.4
	1	89.9	3.9	3.9	4.2	3.9
	10	89.5	6.2	4.3	5.7	4.2

Table 1 also displays results of the approximate variance  $Var\left[\sum_i \hat{r}_i(t)\right]$  obtained using (4.10) and (4.12) and solving the differential equations (2.8) and the second moment equations in Appendix A, (A.27) – (A.32). The equations were solved numerically using the 4<sup>th</sup>/5<sup>th</sup> order Runge-Kutta-Fehlberg method as implemented in MATLAB; (cf. Math Works, 1992).

Table 1 also displays the simple Poisson approximation to  $Var\left[\sum_i \hat{r}_i(t)\right]$  of (5.7) with the  $\alpha_i$  in (3.6).

Recall that the estimate  $Var\left[\sum_i \hat{r}_i(t)\right]$  is unbiased for both approximations. The true value of  $\sum r_i$  is 90. The differential equation approximation is close to the square root of the sample variance for all values of  $\gamma$  for times 10, 20, and 50. The Poisson approximation is close to the square root of the sum of the estimated approximate variances; both of these approximations are neglecting the covariances induced by the purposeful allocation; these covariances become more pronounced as  $\gamma$  becomes larger. There is no covariance for  $\gamma=0$ , equally likely allocation. Note that the Poisson approximation is conservative. However for  $\gamma=1$ , the Poisson approximation is within about 10% of the differential equation approximation which incorporates the covariances. However the difference is larger for  $\gamma=10$ .

Note that the square root of the sample variance and the differential equation approximation suggest that purposeful allocation with  $\gamma=1$  yields the smallest variance of the estimated sum of the numbers of units on all the nodes. A rationale for this suggestion follows.

Suppose there are a fixed number of looks  $K$  that the sensor can take of all nodes and the number of units on each node  $i$ ,  $r_i$ , is known along with the

probability of detecting a unit on node  $i$ ,  $p_i$ . Let  $k_i$  be the number of looks the sensor gives to node  $i$ . If each observation has a binomial distribution

$$\begin{aligned} \text{Var} \left[ \sum_{i=1}^3 \hat{r}_i(t) \right] &= \sum_{i=1}^3 \frac{k_i r_i p_i (1-p_i)}{k_i^2 p_i^2} \\ &= \sum_{i=1}^3 \frac{r_i (1-p_i)}{k_i p_i}. \end{aligned}$$

Lagrange multipliers can be used to show that the (approximate)  $k_i$ ,  $i = 1, 2, 3$  that minimize  $\text{Var} \left[ \sum_{i=1}^3 \hat{r}_i(t) \right]$  are  $k_i = [r_i(1-p_i)/p_i]^{\frac{1}{2}}$ . This solution corresponds to the  $\alpha_i$  of (3.6) with  $a_i = 0$  and  $\gamma = 1$ . Thus, if one is interested in minimizing the estimated variance of the sum of the number of units on all the nodes, then one should look at node  $i$  a number of times proportional to  $[r_i(1-p_i)/p_i]^{0.5}$ . If one were interested in minimizing the estimated variance of the estimate of the number of units on the node with the greatest number of units then one would allocate all looks to that node; this corresponds to the purposeful allocation policy of  $\gamma = \infty$ .

Tables 2 and 3 present results of the simulation experiment with  $r_1 = 49$ ,  $r_2 = 25$ ,  $r_3 = 16$  and  $p_1 = 0.7$ ,  $p_2 = 0.8$ , and  $p_3 = 0.9$ . Table 2 presents the simulation and approximation results for the estimate of the sum of units on all the nodes. Table 3 presents the simulation and approximation results for the number of units on the individual nodes. The differential equation results are close to the simulated values for times  $t = 10, 20, 50$ . The Poisson approximation also seems to be adequate. The Poisson approximation may be doing better in this case because the probabilities of unit detection are larger. One source of the covariance between the estimators  $\hat{r}_i(t)$  is the possibility that  $V(i, t)$  may be 0, in



which case node  $i$  will not be visited very frequently for  $\gamma > 0$  for the purposeful allocation with function  $h_i$  as in (3.1).

**Table 2**  
**Estimate of Total Number of Units on All Nodes**  
 $r_1 = 49, r_2 = 25, r_3 = 16; p_1 = 0.7, p_2 = 0.8, p_3 = 0.9$

		Simulation			Approximation	
		$\sum_i \hat{r}_i(t)$		$\sqrt{\sum_i \sigma_i^2(t)}$	Square Root of Differential Equation Approx Variance	Square Root of Poisson Approx Variance
Time	$\gamma$	Sample Mean	Square Root of Sample Variance	Sample Mean		
5	0	89.9	5.52	3.70	3.30	3.30
	1	89.4	6.53	3.15	3.11	3.02
	10	88.9	8.55	3.14	3.06	2.98
10	0	90.1	2.75	2.84	2.59	2.59
	1	89.9	2.37	2.45	2.46	2.35
	10	89.7	2.46	2.53	2.51	2.35
20	0	90.1	2.04	2.07	1.95	1.95
	1	90.0	1.72	1.80	1.90	1.76
	10	89.9	1.87	1.88	2.06	1.80
50	0	90.0	1.30	1.32	1.28	1.28
	1	90.0	1.12	1.17	1.29	1.16
	10	89.9	1.26	1.24	1.35	1.21



**Table 3**  
Simulation: 500 replications  
Estimate of Number of Units on Node

Time	$\gamma$	Node 1			Node 2			Node 3		
		Mean	Square Root Sample Variance (Diff Eqtn Approx)	$\sqrt{\hat{\sigma}_1^2(t)}$	Mean	Square Root Sample Variance (Diff Eqtn Approx)	$\sqrt{\hat{\sigma}_1^2(t)}$	Mean	Square Root Sample Variance (Diff Eqtn Approx)	$\sqrt{\hat{\sigma}_1^2(t)}$
5	0	49.0	3.89 (2.81)	3.11 (2.81)	24.9	2.20 (1.53)	1.68 (1.53)	16.0	1.17 (0.82)	0.87 (0.82)
	1	48.6	3.95 (2.27)	2.37 (2.38)	24.8	2.38 (1.62)	1.69 (1.59)	16.0	1.54 (1.07)	1.13 (0.99)
	10	48.5	4.85 (2.00)	2.12 (2.17)	24.6	2.94 (1.88)	1.89 (1.70)	15.8	1.97 (1.33)	1.32 (1.13)
10	0	49.1	2.33 (2.20)	2.39 (2.20)	25.0	1.36 (1.20)	1.30 (1.20)	16.0	0.71 (0.64)	0.68 (0.64)
	1	49.0	1.81 (1.74)	1.81 (1.81)	24.9	1.36 (1.27)	1.32 (1.25)	16.0	0.96 (0.89)	0.93 (0.82)
	10	48.9	1.57 (1.53)	1.61 (1.63)	24.9	1.50 (1.44)	1.48 (1.37)	15.9	1.33 (1.29)	1.26 (1.00)
20	0	49.1	1.68 (1.66)	1.75 (1.66)	25.0	0.98 (0.90)	0.95 (0.90)	16.0	0.52 (0.48)	0.50 (0.48)
	1	49.0	1.32 (1.30)	1.32 (1.33)	24.9	0.97 (0.96)	0.98 (0.95)	16.0	0.71 (0.68)	0.71 (0.65)
	10	49.0	1.16 (1.15)	1.19 (1.19)	24.9	1.08 (1.08)	1.10 (1.05)	15.9	1.06 (1.00)	0.95 (0.84)
50	0	49.1	1.08 (1.09)	1.12 (1.09)	25.0	0.63 (0.59)	0.61 (0.59)	16.0	0.33 (0.32)	0.32 (0.32)
	1	49.0	0.88 (0.85)	0.86 (0.86)	25.0	0.64 (0.63)	0.64 (0.63)	16.0	0.45 (0.45)	0.46 (0.44)
	10	49.0	0.77 (0.76)	0.77 (0.77)	24.9	0.72 (0.72)	0.72 (0.70)	16.0	0.71 (0.66)	0.65 (0.61)

## REFERENCES

- P. J. Bickel and K. A. Doksum, "*Mathematical Statistics: Basic Ideas and Selected Topics*," Holden-Day, Inc., San Francisco, CA, 1977.
- D. P. Gaver and P. A. Jacobs, "Asymptotic properties of stochastic greedy bin-packing," *Stochastic Models*, **11** (1995) pp. 333-348.
- D. P. Gaver, J. A. Morrison, and R. Silveira, "Service-adaptive multitype repairman problems," *SIAM J. Appl. Math.*, **53** (1993) pp. 454-470.
- P. Hall, "On the erratic behavior of estimators of  $N$  in the binomial  $N, p$  distribution," *Journal of the American Statistical Association*, **89** (1994) pp. 344-352.
- The Math Works, Inc., *MATLAB Reference Guide*, The Math Works, Inc., Natick, MA, August 1992.
- S. K. Thompson, and G. A. F. Seber, "Detectability in conventional and adaptive sampling," *Biometrics*, **50** (1994) pp. 712-725.

## APPENDIX A

In the Appendix we present details of the normal approximation. We follow an analytical approach used in a different context in Gaver and Jacobs (1995), and in Gaver, Morrison, and Silveira (1993). Let the moment-generating function (assumed to exist, otherwise use the characteristic function) be

$$\begin{aligned}\psi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) &= E[\exp\{\boldsymbol{\theta}N(t) + \boldsymbol{\xi}V(t)\}] \\ &= E\left[\exp\left\{\sum_{j=1}^I \theta_j N_j(t) + \sum_{j=1}^I \xi_j V_j(t)\right\}\right].\end{aligned}\tag{A.1}$$

Condition on  $(N_i(t), V_i(t))$ ,  $i \in \{1, 2, \dots, I\}$  to obtain

$$\begin{aligned}& E\left[\exp\left\{\sum_{j=1}^I \theta_j N_j(t+h) + \sum_{j=1}^I \xi_j V_j(t+h)\right\} \middle| N(t), V(t)\right] \\ &= (1 - \lambda h) \exp\{\boldsymbol{\theta}N(t) + \boldsymbol{\xi}V(t)\} \\ & \quad + \lambda h \sum_i \alpha_i(N(t), V(t)) \exp\{\boldsymbol{\theta}N(t) + \boldsymbol{\xi}V(t)\} [e^{\theta_i \hat{b}(\xi_i)}]\end{aligned}\tag{A.2}$$

where

$$\hat{b}(\xi_i) = E[e^{\xi_i Z(i)}].\tag{A.3}$$

with  $Z(i)$  an observation of the number of units on node  $i$ .

Let  $h \rightarrow 0$  to obtain

$$\frac{\partial}{\partial t} \psi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) = -\lambda \psi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) + \lambda \sum_{i=1}^I E[\alpha_i(N(t), V(t)) \exp\{\boldsymbol{\theta}N(t) + \boldsymbol{\xi}V(t)\} e^{\theta_i \hat{b}(\xi_i)}].\tag{A.4}$$

### Scaling

Let

$$X_j(t) = \frac{N_j(t) - \lambda m_j(t)}{\sqrt{\lambda}}\tag{A.5}$$

$$Y_j(t) = \frac{V_j(t) - \lambda w_j(t)}{\sqrt{\lambda}} \quad (\text{A.6})$$

and let  $\lambda \gg 1$ .

Let

$$\varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) = E[\exp\{\boldsymbol{\theta}X(t) + \boldsymbol{\xi}Y(t)\}]; \quad (\text{A.7})$$

then

$$\psi(\boldsymbol{\theta}/\sqrt{\lambda}, \boldsymbol{\xi}/\sqrt{\lambda}; t) = \varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \exp\{\sqrt{\lambda}[\boldsymbol{\theta}m(t) + \boldsymbol{\xi}v(t)]\}. \quad (\text{A.8})$$

Thus, we have the following equation from (A.8) and (A.4)

$$\begin{aligned} & \frac{\partial}{\partial t} \psi(\boldsymbol{\theta}/\sqrt{\lambda}, \boldsymbol{\xi}/\sqrt{\lambda}; t) \\ &= \frac{\partial}{\partial t} \varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \exp\{\sqrt{\lambda}[\boldsymbol{\theta}m(t) + \boldsymbol{\xi}v(t)]\} \\ &+ \sqrt{\lambda} \varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \exp\{\sqrt{\lambda}[\boldsymbol{\theta}m(t) + \boldsymbol{\xi}v(t)]\} [\boldsymbol{\theta}m'(t) + \boldsymbol{\xi}v'(t)] \\ &= \left[ -\lambda \varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \right. \end{aligned} \quad (\text{A.9})$$

$$\begin{aligned} & \left. + \lambda \sum_{i=1}^I E \left[ \alpha_i \left( m(t) + \frac{1}{\sqrt{\lambda}} X(t), v(t) + \frac{1}{\sqrt{\lambda}} Y(t) \right) \exp\{\boldsymbol{\theta}X(t) + \boldsymbol{\xi}Y(t)\} e^{\theta_i/\sqrt{\lambda} \hat{b}_i(\xi_i/\sqrt{\lambda})} \right] \right] \\ & \times \exp\{\sqrt{\lambda}[\boldsymbol{\theta}m(t) + \boldsymbol{\xi}v(t)]\}. \end{aligned}$$

Dividing both sides by  $\exp\{\sqrt{\lambda}[\boldsymbol{\theta}m(t) + \boldsymbol{\xi}v(t)]\}$  we obtain

$$\begin{aligned} & \frac{\partial}{\partial t} \varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) + \sqrt{\lambda} \varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) [\boldsymbol{\theta}m'(t) + \boldsymbol{\xi}v'(t)] \\ &= -\lambda \varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \\ &+ \lambda \sum_{i=1}^I E \left[ \left\{ \alpha_i(m(t), v(t)) + \sum_j \frac{\partial}{\partial m_j} \alpha_i(m(t), v(t)) \frac{1}{\sqrt{\lambda}} X_j(t) \right. \right. \\ & \left. \left. + \sum_j \frac{\partial}{\partial \beta_j} \alpha_i(m(t), v(t)) \frac{1}{\sqrt{\lambda}} Y_j(t) + O\left(\frac{1}{\sqrt{\lambda}}\right) \right\} \exp\{\boldsymbol{\theta}X(t) + \boldsymbol{\xi}Y(t)\} e^{\theta_i/\sqrt{\lambda} \hat{b}_i(\xi_i/\sqrt{\lambda})} \right] \end{aligned} \quad (\text{A.10})$$

Let

$$\alpha_i(m(t), v(t)) = \frac{h_i(m_i(t), v_i(t))}{\sum_j h_j(m_j(t), v_j(t))} \quad (\text{A.11})$$

where  $h_j(x, y)$  is a sufficiently smooth function with first order partial derivatives and  $h_j(\lambda^p x, \lambda^p y) = \lambda^p h_j(x, y)$ .

Let

$$H_i(x; t) = \frac{\frac{\partial}{\partial m_i} h_i(m_i(t), v_i(t))}{\sum_j h_j(m_j(t), v_j(t))} \quad (\text{A.12})$$

$$H_i(y; t) = \frac{\frac{\partial}{\partial v_i} h_i(m_i(t), v_i(t))}{\sum_j h_j(m_j(t), v_j(t))} \quad (\text{A.13})$$

$$H_{ik}(x; t) = \frac{h_i(m_i(t), v_i(t))}{\left( \sum_j h_j(m_j(t), v_j(t)) \right)^2} \frac{\partial}{\partial m_k} h_k(m_k(t), v_k(t)) \quad (\text{A.14})$$

$$H_{ik}(y; t) = \frac{h_i(m_i(t), v_i(t))}{\left( \sum_j h_j(m_j(t), v_j(t)) \right)^2} \frac{\partial}{\partial v_k} h_k(m_k(t), v_k(t)) \quad (\text{A.15})$$

Note that

$$\begin{aligned} & \alpha_i(\lambda m_i(t) + \sqrt{\lambda} X_i(t), \lambda v_i(t) + \sqrt{\lambda} Y_i(t)) \\ &= \frac{h_i(m_i(t), v_i(t))}{\sum_{j=1}^I h_j(m_j(t), v_j(t))} \end{aligned} \quad (\text{A.16})$$

$$+ \frac{1}{\sqrt{\lambda}} \left[ H_i(x; t) X_i(t) + H_i(y; t) Y_i(t) + \sum_k H_{ik}(x; t) X_k(t) + \sum_k H_{ik}(y; t) Y_k(t) \right] + O\left(\frac{1}{\lambda}\right).$$

Since

$$\sum_i \alpha_i (\lambda m_i(t) + \sqrt{\lambda} X_i(t), \lambda v_i(t) + \sqrt{\lambda} Y_i(t)) = 1 \quad (\text{A.17})$$

this implies that the summed coefficients of  $1/\sqrt{\lambda}$ ,  $1/\lambda$  etc. must individually be 0.

Expression (A.10) can be rewritten as

$$\begin{aligned} & \frac{\partial}{\partial t} \varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) + \sqrt{\lambda} \varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) [\boldsymbol{\theta} \mathbf{m}'(t) + \boldsymbol{\xi} \mathbf{v}'(t)] \\ &= -\lambda \varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \\ &+ \lambda \sum_i \left[ 1 + \frac{\theta_i}{\sqrt{\lambda}} + \frac{b_1(i) \xi_i}{\sqrt{\lambda}} + \frac{1}{2} \frac{\theta_i^2}{\lambda} + \frac{1}{2} \frac{b_2(i) \xi_i^2}{\lambda} \right] \\ & \times \left\{ \frac{h_i(m_i(t), v_i(t))}{\sum_j h_j(m_j(t), v_j(t))} \varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \right. \\ &+ \frac{1}{\sqrt{\lambda}} \left[ H_i(x; t) \frac{\partial}{\partial \theta_i} \varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) + H_i(y; t) \frac{\partial}{\partial \xi_i} \varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \right. \\ & \left. \left. - \sum_{k=1}^I \left[ H_{ik}(x; t) \frac{\partial}{\partial \theta_k} \varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) + H_{ik}(y; t) \frac{\partial}{\partial \xi_k} \varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \right] \right] \right\} + O\left(\frac{1}{\lambda}\right) \end{aligned} \quad (\text{A.18})$$

where  $b_n(i) = E[Z(i)^n]$  the  $n^{\text{th}}$  moment of an observation at node  $i$ .

Let

$$\varphi(\boldsymbol{\theta}, \boldsymbol{\xi}; t) = \sum_{\ell=0}^{\infty} \varphi_{\ell}(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \lambda^{-\ell/2}. \quad (\text{A.19})$$

Substituting (A.19) into (A.18) results in the following equation for  $\varphi_0$



$$\begin{aligned}
& \frac{\partial}{\partial t} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) + \sqrt{\lambda} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) [\boldsymbol{\theta} m'(t) + \boldsymbol{\xi} v'(t)] \\
& = -\lambda \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \\
& + \lambda \sum_i \left[ 1 + \frac{1}{\sqrt{\lambda}} (\theta_i + b_1(i) \xi_i) + \frac{1}{2} \frac{1}{\lambda} (\theta_i^2 + b_2(i) \xi_i^2) \right] \\
& \times \left\{ \frac{h_i(m_i(t), v_i(t))}{\sum_j h_j(m_j(t), v_j(t))} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \right. \\
& + \frac{1}{\sqrt{\lambda}} \left[ H_i(x; t) \frac{\partial}{\partial \theta_i} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) + H_i(y; t) \frac{\partial}{\partial \xi_i} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \right. \\
& \left. \left. - \sum_{k=1}^I \left[ H_{ik}(x; t) \frac{\partial}{\partial \theta_k} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) + H_{ik}(y; t) \frac{\partial}{\partial \xi_k} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \right] \right] \right\} + O\left(\frac{1}{\lambda}\right). \tag{A.20}
\end{aligned}$$

Equating terms of order  $\lambda^{l/2}$ , the terms of order  $\lambda$  cancel. The terms of order  $\sqrt{\lambda}$  result in the equation

$$\begin{aligned}
& \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) [\boldsymbol{\theta} m'(t) + \boldsymbol{\xi} v'(t)] \\
& = \sum_{i=1}^I \left\{ \frac{h_i(m_i(t), v_i(t))}{\sum_j h_j(m_j(t), v_j(t))} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) [\theta_i + b_1(i) \xi_i] \right. \\
& + H_i(x; t) \frac{\partial}{\partial \theta_i} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) + H_i(y; t) \frac{\partial}{\partial \xi_i} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \\
& \left. + \sum_{k=1}^I \left[ H_{ik}(x; t) \frac{\partial}{\partial \theta_k} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) + H_{ik}(y; t) \frac{\partial}{\partial \xi_k} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \right] \right\} + O\left(\frac{1}{\lambda}\right). \tag{A.21}
\end{aligned}$$

The terms of order  $\sqrt{\lambda}$  cancel if

$$\boldsymbol{\theta}m'(t) + \boldsymbol{\xi}v'(t) = \sum_{i=1}^I \frac{h_i(m_i(t), v_i(t))}{\sum_j h_j(m_j(t), v_j(t))} [\theta_i + b_1(i)\xi_i]. \quad (\text{A.22})$$

In order for this to occur

$$\frac{d}{dt} m_i(t) = \frac{h_i(m_i(t), v_i(t))}{\sum_j h_j(m_j(t), v_j(t))} \quad i = 1, \dots, I \quad (\text{A.23})$$

and

$$\frac{d}{dt} v_i(t) = b_1(i) \frac{h_i(m_i(t), v_i(t))}{\sum_j h_j(m_j(t), v_j(t))} \quad i = 1, \dots, I. \quad (\text{A.24})$$

Thus,

$$v_i(t) = b_1(i)m_i(t). \quad (\text{A.25})$$

Next look for terms of order 1 in (A.20).

$$\begin{aligned} & \frac{\partial}{\partial t} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \\ &= \sum_i \frac{h_i(m_i(t), v_i(t))}{\sum_j h_j(m_j(t), v_j(t))} \left[ \frac{1}{2} \theta_i^2 + \frac{1}{2} b_2(i) \xi_i^2 + b_1(i) \theta_i \xi_i \right] \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \\ &+ \sum_i \left[ H_i(x; t) \frac{\partial}{\partial \theta_i} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) + H_i(y; t) \frac{\partial}{\partial \xi_i} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \right] [\theta_i + b_1(i) \xi_i] \\ &+ \sum_i \sum_k \left[ H_{ik}(x; t) \frac{\partial}{\partial \theta_k} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) + H_{ik}(y; t) \frac{\partial}{\partial \xi_k} \varphi_0(\boldsymbol{\theta}, \boldsymbol{\xi}; t) \right] [\theta_i + b_1(i) \xi_i]. \end{aligned} \quad (\text{A.26})$$

Equations for the joint moments of  $\{(X_j(t), Y_j(t))\}$  can be obtained by differentiating (A.26) with respect to  $\{\theta_i\}$  and  $\{\xi_i\}$  and evaluated at  $\boldsymbol{\theta} = \boldsymbol{\xi} = 0$ . The resulting equations are

$$\begin{aligned}
& \frac{d}{dt} E[X_\ell^2(t)] \\
&= \frac{h(m_\ell(t), v_\ell(t))}{\sum_j h(m_j(t), v_j(t))} + 2H_\ell(x; t) E[X_\ell(t)^2] + 2H_\ell(y; t) E[X_\ell(t)Y_\ell(t)] \\
&+ 2\sum_k H_{\ell k}(x; t) E[X_\ell(t)X_k(t)] + 2\sum_k H_{\ell k}(y; t) E[X_\ell(t)Y_k(t)]
\end{aligned} \tag{A.27}$$

For  $\ell \neq a$

$$\begin{aligned}
& \frac{d}{dt} E[X_\ell(t)X_a(t)] \\
&= H_a(x; t) E[X_a(t)X_\ell(t)] + H_a(y; t) E[Y_a(t)X_\ell(t)] \\
&+ H_\ell(x; t) E[X_a(t)X_\ell(t)] + H_\ell(y; t) E[X_a(t)Y_\ell(t)] \\
&+ \sum_k H_{ak}(x; t) E[X_k(t)X_\ell(t)] + \sum_k H_{ak}(y; t) E[X_\ell(t)Y_k(t)] \\
&+ \sum_k H_{\ell k}(x; t) E[X_a(t)X_k(t)] + \sum_k H_{\ell k}(y; t) E[X_a(t)Y_k(t)].
\end{aligned} \tag{A.28}$$

$$\begin{aligned}
\frac{d}{dt} E[Y_\ell^2(t)] &= \frac{h(m_\ell(t), v_\ell(t))}{\sum_j h(m_j(t), v_j(t))} b_2(\ell) \\
&+ 2b_1(\ell) \left[ H_\ell(x; t) E[X_\ell(t)Y_\ell(t)] + H_\ell(y; t) E[Y_\ell^2(t)] \right] \\
&+ 2b_1(\ell) \sum_j H_{\ell j}(x; t) E[X_j(t)Y_\ell(t)] + H_{\ell j}(y; t) E[Y_\ell(t)Y_j(t)]
\end{aligned} \tag{A.29}$$

For  $j \neq k$

$$\begin{aligned}
& \frac{d}{dt} E[Y_k(t)Y_j(t)] \\
&= \left\{ H_j(x;t)E[X_j(t)Y_k(t)] + H_j(y;t)E[Y_j(t)Y_k(t)] \right\} b_1(j) \\
&+ \left\{ H_k(x;t)E[X_k(t)Y_j(t)] + H_k(y;t)E[Y_j(t)Y_k(t)] \right\} b_1(k) \\
&+ b_1(j) \sum_{\ell} H_{j\ell}(x;t)E[X_{\ell}(t)Y_k(t)] + H_{j\ell}(y;t)E[Y_k(t)Y_{\ell}(t)] \\
&+ b_1(k) \sum_{\ell} H_{k\ell}(x;t)E[X_{\ell}(t)Y_j(t)] + H_{k\ell}(y;t)E[Y_{\ell}(t)Y_j(t)]
\end{aligned} \tag{A.30}$$

$$\begin{aligned}
& \frac{d}{dt} E[X_k(t)Y_k(t)] \\
&= \frac{h_k(m_k(t), v_k(t))}{\sum_j h_j(m_j(t), v_j(t))} b_1(k) \\
&+ H_k(x;t) \left\{ E[Y_k(t)X_k(t)] + E[X_k^2(t)] b_1(k) \right\} \\
&+ H_k(y;t) \left\{ E[Y_k^2(t)] + E[X_k(t)Y_k(t)] b_1(k) \right\} \\
&+ \sum_j \left\{ H_{kj}(x;t)E[X_j(t)Y_k(t)] + H_{kj}(y;t)E[Y_k(t)Y_j(t)] \right\} \\
&+ \sum_j \left\{ H_{kj}(x;t)E[X_j(t)X_k(t)] + H_{kj}(y;t)E[X_k(t)Y_j(t)] \right\} b_1(k)
\end{aligned} \tag{A.31}$$

For  $\ell \neq k$

$$\begin{aligned} & \frac{d}{dt} E[X_\ell(t)Y_k(t)] \\ &= H_\ell(x;t)E[X_\ell(t)Y_k(t)] + H_\ell(y;t)E[Y_\ell(t)Y_k(t)] \\ &+ \{H_k(x;t)E[X_\ell(t)X_k(t)] + H_k(y;t)E[X_\ell(t)Y_k(t)]\}b_1(k) \\ &+ \sum_j H_{\ell j}(x;t)E[X_j(t)Y_k(t)] + H_{\ell j}(y;t)E[Y_k(t)Y_j(t)] \\ &+ \sum_j \{H_{kj}(x;t)E[X_j(t)X_\ell(t)] + H_{kj}(y;t)E[Y_j(t)X_\ell(t)]\}b_1(k) \end{aligned} \tag{A.32}$$





## INITIAL DISTRIBUTION LIST

1.	Research Office (Code 08) .....	1
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
2.	Dudley Knox Library (Code 52) .....	2
	Naval Postgraduate School	
	Monterey, CA 93943-5002	
3.	Defense Technical Information Center .....	2
	Cameron Station	
	Alexandria, VA 22314	
4.	Director, Force Structure, Resources and Assessment (J-8) .....	2
	8000 The Joint Staff	
	Washington, DC 20310-8000	
5.	Deputy Director for Technical Operations, J-8 .....	1
	8000 The Joint Staff	
	Washington, DC 20310-8000	
6.	Chief, Warfare Analysis Division, J-8 .....	1
	8000 The Joint Staff	
	Washington, DC 20310-8000	
7.	Prof. Donald P. Gaver (Code OR/Gv) .....	10
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
8.	Prof. Patricia A. Jacobs (Code OR/Jc) .....	5
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
9.	Prof. Mark Youngren (Code OR/Ym) .....	5
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
10.	Prof. George Connor (Code OR/Co) .....	1
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
11.	Prof. Michael Sovereign (Code OR/Sm) .....	1
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
12.	Ms. Therese Bilodeau (Code OR/Bi) .....	1
	Naval Postgraduate School	
	Monterey, CA 93943-5000	

13. Dr. J. Abrahams, Code 1111, Room 607 ..... 1  
 Mathematical Sciences Division, Office of Naval Research  
 800 North Quincy Street  
 Arlington, VA 22217-5000
14. Dr. John Bailer ..... 1  
 468 N St. NW  
 Washington, DC 20024
15. Prof. D. R. Barr ..... 1  
 Dept. of Systems Engineering  
 U.S. Military Academy  
 West Point, NY 10996
16. Dr. David Brillinger ..... 1  
 Statistics Department  
 University of California  
 Berkeley, CA 94720
17. Dr. David Burman ..... 1  
 AT&T Bell Telephone Laboratories  
 600 Mountain Avenue  
 Murray Hill, NJ 07974
18. Prof. Brad Carlin ..... 1  
 School of Public Health  
 University of Minnesota  
 Mayo Bldg. A460  
 Minneapolis, MN 55455
19. Dr. Robert Carpenter ..... 1  
 NAMRI/Navy Toxicology Detachment  
 Bldg. 433, Area B  
 Wright-Patterson AFB, OH 45433-6503
20. Center for Naval Analyses ..... 1  
 4401 Ford Avenue  
 Alexandria, VA 22302-0268
21. Prof. H. Chernoff ..... 1  
 Department of Statistics  
 Harvard University  
 1 Oxford Street  
 Cambridge, MA 02138
22. Mr. Wm. P. Clay ..... 1  
 Director, USAMSAA  
 Attn: AMXSY-CA  
 APG, MD 21005-5071
23. Dr. Edward G. Coffman, Jr. .... 1  
 AT&T Bell Telephone Laboratories  
 600 Mountain Avenue  
 Murray Hill, NJ 07974

24. Dr. John Copas ..... 1  
 Dept. of Mathematics,  
 University of Birmingham  
 P. O. Box 363  
 Birmingham B15 2TT  
 ENGLAND
25. Professor Sir David Cox ..... 1  
 Nuffield College  
 Oxford, OX1 INF  
 ENGLAND
26. Professor H. G. Daellenbach ..... 1  
 Department of Operations Research  
 University of Canterbury  
 Christchurch, NEW ZEALAND
27. Dr. S. R. Dalal ..... 1  
 Bellcore  
 445 South Street  
 Morristown, NJ 07962-1910
28. Dr. D. F. Daley ..... 1  
 Statistic Dept. (I.A.S.)  
 Australian National University  
 Canberra, A.C.T. 2606  
 AUSTRALIA
29. Dr. B. Doshi ..... 1  
 AT&T Bell Laboratories  
 HO 3M-335  
 Holmdel, NJ 07733
30. Dr. Naihua Duan ..... 1  
 RAND Corporation  
 Santa Monica, CA 90406
31. Prof. Bradley Efron ..... 1  
 Statistics Dept.  
 Sequoia Hall  
 Stanford University  
 Stanford, CA 94305
32. Dr. Guy Fayolle ..... 1  
 I.N.R.I.A.  
 Dom de Voluceau-Rocquencourt  
 78150 Le Chesnay Cedex  
 FRANCE
33. Dr. Andrew Gelman ..... 1  
 Statistics Dept.  
 University of California  
 Berkeley, CA 94720

34.	Dr. Neil Gerr .....	1
	Office of Naval Research Arlington, VA 22217	
35.	Prof. Peter Glynn .....	1
	Dept. of Operations Research Stanford University Stanford, CA 94350	
36.	Prof. Bernard Harris.....	1
	Dept. of Statistics University of Wisconsin 610 Walnut Street Madison, WI 53706	
37.	Prof. J. Michael Harrison.....	1
	Graduate School of Business Stanford University Stanford, CA 94305-5015	
38.	Dr. P. Heidelberger .....	1
	IBM Research Laboratory Yorktown Heights New York, NY 10598	
39.	Dr. D. C. Hoaglin .....	1
	Department of Statistics Harvard University 1 Oxford Street Cambridge, MA 02138	
40.	Prof. D. L. Iglehart .....	1
	Dept. of Operations Research Stanford University Stanford, CA 94350	
41.	Institute for Defense Analysis .....	1
	1800 North Beauregard Alexandria, VA 22311	
42.	Prof. J. B. Kadane .....	1
	Dept. of Statistics Carnegie-Mellon University Pittsburgh, PA 15213	
43.	Dr. F. P. Kelly .....	1
	Statistics Laboratory 16 Mill Lane Cambridge ENGLAND	

44. Dr. Jon Kettenring ..... 1  
 Bellcore  
 445 South Street  
 Morris Township, NJ 07962-1910
45. Koh Peng Kong ..... 1  
 OA Branch, DSO  
 Ministry of Defense  
 Blk 29 Middlesex Road  
 SINGAPORE 1024
46. Prof. Guy Latouche ..... 1  
 University Libre Bruxelles  
 C.P. 212, Blvd. De Triomphe  
 B-1050 Bruxelles  
 BELGIUM
47. Dr. A. J. Lawrance ..... 1  
 Dept. of Mathematics,  
 University of Birmingham  
 P. O. Box 363  
 Birmingham B15 2TT  
 ENGLAND
48. Prof. M. Leadbetter ..... 1  
 Department of Statistics  
 University of North Carolina  
 Chapel Hill, NC 27514
49. Prof. J. Lehoczky ..... 1  
 Department of Statistics  
 Carnegie-Mellon University  
 Pittsburgh, PA 15213
50. Dr. James R. Maar ..... 1  
 National Security Agency  
 9608 Basket Ring  
 Columbia, MD 21045-0689
51. Dr. Colin Mallows ..... 1  
 AT&T Bell Telephone Laboratories  
 600 Mountain Avenue  
 Murray Hill, NJ 07974
52. Prof. R. Douglas Martin ..... 1  
 Department of Statistics, GN-22  
 University of Washington  
 Seattle, WA 98195
53. Prof. M. Mazumdar ..... 1  
 Dept. of Industrial Engineering  
 University of Pittsburgh  
 Pittsburgh, PA 15235

54.	Dr. James McKenna .....	1
	Bell Communications Research	
	445 South Street	
	Morristown, NJ 07960-1910	
55.	Prof. Paul Moose .....	1
	C3I Academic Group	
	Naval Postgraduate School	
	Monterey, CA 93943-5000	
56.	Prof. Carl N. Morris.....	1
	Statistics Department	
	Harvard University	
	1 Oxford St.	
	Cambridge, MA 02138	
57.	Dr. John A. Morrison .....	1
	AT&T Bell Telephone Laboratories	
	600 Mountain Avenue	
	Murray Hill, NJ 07974	
58.	Prof. F. W. Mosteller .....	1
	Department of Statistics	
	Harvard University	
	1 Oxford St.	
	Cambridge, MA 02138	
59.	Operations Research Center, Rm E40-164 .....	1
	Massachusetts Institute of Technology	
	Attn: R. C. Larson and J. F. Shapiro	
	Cambridge, MA 02139	
60.	Dr. T. J. Ott .....	1
	Bellcore,	
	445 South Street	
	Morristown, NJ 07962-1910	
	(MRE 2P388)	
61.	Dr. Jim Petty .....	1
	National Biological Survey	
	4200 New Haven Road	
	Columbia, MO 65201	
62.	Dr. V. Ramaswami .....	1
	MRE 2Q-358	
	Bell Communications Research, Inc.	
	445 South Street	
	Morristown, NJ 07960	
63.	Dr. Martin Reiman .....	1
	Rm #2C-117	
	AT&T Bell labs	
	600 Mountain Ave.	
	Murray Hill, NJ 07974-2040	



64. Prof. Maria Rieders ..... 1  
 Dept. of Industrial Eng.  
 Northwestern Univ.  
 Evanston, IL 60208
65. Dr. Rhonda Righter ..... 1  
 Dept. of Decision & Info. Sciences  
 Santa Clara University  
 Santa Clara, CA 95118
66. Dr. John E. Rolph ..... 1  
 RAND Corporation  
 1700 Main St.  
 Santa Monica, CA 90406
67. Prof. Frank Samaniego ..... 1  
 Statistics Department  
 University of California  
 Davis, CA 95616
68. Prof. W. R. Schucany ..... 1  
 Dept. of Statistics  
 Southern Methodist University  
 Dallas, TX 75222
69. Prof. G. A. F. Seber ..... 1  
 Dept. of Statistics  
 Univ. of Auckland  
 Private Bag 92019  
 Auckland, NEW ZEALAND
70. Prof. G. Shantikumar ..... 1  
 The Management Science Group  
 School of Business Administration  
 University of California  
 Berkeley, CA 94720
71. Prof. D. C. Siegmund ..... 1  
 Dept. of Statistics  
 Sequoia Hall  
 Stanford University  
 Stanford, CA 94305
72. Prof. N. D. Singpurwalla ..... 1  
 George Washington University  
 Washington, DC 20052
73. Prof. H. Solomon ..... 1  
 Department of Statistics  
 Sequoia Hall  
 Stanford University  
 Stanford, CA 94305

74.	Dr. Andrew Solow .....	1
	Woods Hole Oceanographic Institute Woods Hole, MA 02543	
75.	Prof. W. Stuetzle .....	1
	Department of Statistics University of Washington Seattle, WA 98195	
76.	Prof. J. R. Thompson .....	1
	Dept. of Mathematical Science Rice University Houston, TX 77001	
77.	Prof. Steven K. Thompson .....	1
	Statistics Dept. Pennsylvania State Univ. 326 Classroom Bldg. University Park, PA 16802-2111	
78.	Prof. J. W. Tukey .....	1
	Statistics Dept., Fine Hall Princeton University Princeton, NJ 08540	
79.	Dr. D. Vere-Jones .....	1
	Dept. of Math, Victoria Univ. of Wellington P. O. Box 196 Wellington NEW ZEALAND	
80.	Prof. David L. Wallace .....	1
	Statistics Dept., University of Chicago 5734 S. University Ave. Chicago, IL 60637	
81.	Dr. Ed Wegman .....	1
	George Mason University Fairfax, VA 22030	
82.	Dr. Alan Weiss .....	1
	Rm. 2C-118 AT&T Bell Laboratories 600 Mountain Avenue Murray Hill, NJ 07974-2040	
83.	Dr. P. Welch .....	1
	IBM Research Laboratory Yorktown Heights, NY 10598	
84.	Prof. Roy Welsch .....	1
	Sloan School M.I.T. Cambridge, MA 02139	



DUDLEY KNOX LIBRARY



3 2768 00326204 9